



# Introduction to Differential Equations Without the Agonizing Pain : Practice Solutions

Makiror Ouyang

## I. First-order ordinary differential equations

(1)  $\frac{dy}{dx} = 2x$

- Tip: Separate the variables and integrate both sides to solve the differential equation.

(2)  $\frac{dy}{dx} + 2y = e^{-x}$

- Tip: Rewrite this in standard form, find appropriate integration factors, express the left-hand side as a differential, and integrate both sides of the equation.

(3)  $\frac{dy}{dx} + (x - 2y) = 0$

- Tip: Check if it is an exact equation. If not, find the integrating factor and convert it into the exact equation.

## Solutions

(1)

We can separate variables and integrate both sides with respect to their respective variables.

$$\begin{aligned}\frac{dy}{dx} &= 2x \\ \int dy &= \int 2x \, dx \\ \int dy &= \frac{1}{2} \cdot 2x^{(1+1)} + C \\ y &= x^2 + C\end{aligned}$$

(2)

First, we can rewrite in Standard Form  $\frac{dy}{dx} + P(x) = Q(x)$ , where  $P(x) = 2$ , and find the potential function  $\phi(x) = e^{\int P(x)dx} = e^{\int 2dx}$

$$\begin{aligned}\frac{dy}{dx} + 2y &= e^{-x} \\ e^{\int 2dx} \left( \frac{dy}{dx} + 2y \right) &= e^{\int 2dx} e^{-x} \\ e^{\int 2dx} \frac{dy}{dx} + e^{\int 2dx} 2y &= 1\end{aligned}$$

Next, we can express the left-hand side as differential:

$$\frac{d}{dx} (e^{2x} y) dx = e^{-x}$$

Finally, integrate both sides and solve for  $y$ :

$$\begin{aligned}\frac{d}{dx} (e^{2x} y) dx &= 1 \\ \int \frac{d}{dx} (e^{2x} y) dx &= \int 1 dx \\ e^{2x} y &= e^{-x} + C \\ y &= e^{-x} + C e^{-2x}\end{aligned}$$

(3)

First we rewrite the equation:

$$\begin{aligned}\frac{dy}{dx} + (x - 2y) &= 0 \\ \frac{dy}{dx} - 2y &= -x\end{aligned}$$

The equation is exact if the partial derivatives with respect to  $y$  of the coefficient of  $dx$  and with respect to  $x$  of the coefficient of  $dy$  are equal.

$$M_x = \frac{\partial}{\partial x} 1 = 0$$

$$N_y = \frac{\partial}{\partial y} -2y = -2$$

$$0 \neq -2$$

Since  $M_x \neq N_y$ , the equation is not exact. The integrating factor  $\mu$  can be found using  $\mu = e^{\int \frac{N_y - M_x}{M} dx}$ . In this case:

$$\mu = e^{\int \frac{N_y - M_x}{M} dx} = e^{\int -2 dx} = e^{-2x}$$

Multiply the entire equation by  $\mu$ :

$$\mu M(x, y) dx + \mu N(x, y) dy = -x$$

$$e^{-2x} \frac{dy}{dx} - 2e^{-2x} y = -e^{-2x} x$$

$$e^{-2x} \frac{dy}{dx} - \frac{d}{dx}(e^{-2x})y = -e^{-2x} x$$

$$\frac{d}{dx}(e^{-2x} y) = -e^{-2x} x$$

$$y = \frac{x}{2} + \frac{1}{4} + Ce^{2x}$$

## II. Linear differential equations with constant coefficients

- (1) Find the roots of the characteristic equation for  $y'' + 2y' + 2y = 0$  and write down its solution.
- (2) Using the Exponential-input Theorem to solve  $\frac{dy}{dx} + y = e^x$ .
- (3) Expand the Laplace transform of the function  $f(x) = e^{2x}$  over the interval  $[0, \infty]$ .
- (4) Find the inverse function of the Laplace transform for the function  $F(s) = \frac{1}{s^2 + 4s + 5}$ .

### Solutions

(1)

To find the roots of the characteristic equation for the given second-order linear homogeneous

differential equation  $y'' + 2y' + 2y = 0$ , we can write down the characteristic equation by replacing the derivatives with the corresponding terms.

The characteristic equation is obtained by substituting  $y = e^{rt}$  into the differential equation:

$$r^2 + 2r + 2 = 0$$

For the given equation,  $a = 1$ ,  $b = 2$ ,  $c = 2$ . The solutions can be found using the quadratic formula:

$$\Delta = b^2 - 4ac = 2^2 - 4 \cdot 1 \cdot 2 = -4$$

$$r = \frac{-b \pm \sqrt{\Delta}}{2a} = \frac{-2 \pm \sqrt{-4}}{2}$$

Since the  $\Delta$  is negative, the roots will be complex conjugates:

$$r = -1 \pm i$$

That's  $\alpha = -1$ ,  $\beta = 1$ . According to Euler's formula  $e^{ix} = \cos(x) + i \sin(x)$ , we can substitute:

$$e^{(\alpha+i\beta)x} = e^{\alpha x} (\cos(\beta x) \pm i \sin(\beta x))$$

So we got the general solution formula:

$$\begin{aligned} y &= C_1 e^{r_1 x} + C_2 e^{r_2 x} \\ &= C_1 e^{\alpha x} (\cos(\beta x) + i \sin(\beta x)) + C_2 e^{\alpha x} (\cos(\beta x) - i \sin(\beta x)) \end{aligned}$$

Finally, by substituting, we got the solution of the equation:

$$y = e^{-x} (C_1 \sin(x) + C_2 \cos(x))$$

(2)

We can rewrite in Standard Form  $\frac{dy}{dx} + P(x)y = Q(x)$ , where  $P(x)$ ,  $Q(x)$  are function of  $x$ , and find the potential function  $\phi(x) = e^{\int P(x)dx} = e^{\int 1dx} = e^x$

$$\begin{aligned}
e^x \frac{dy}{dx} + e^x y &= e^{2x} \\
\frac{d}{dx}(e^x y) &= e^{2x} \\
\int \frac{d}{dx}(e^x y) dx &= \int e^{2x} dx \\
e^x y &= \frac{1}{2} e^{2x} + C \\
y &= \frac{1}{2} e^{2x} + C e^{-x}
\end{aligned}$$

(3)

The Laplace transform of a function  $f(x)$  defined on  $[0, \infty)$  is given by the integral:

$$\mathcal{L}\{f(x)\} = \int_0^{\infty} f(x) e^{-sx} dx$$

In the case:

$$\begin{aligned}
\mathcal{L}\{f(x)\} &= \int_0^{\infty} e^{2x} e^{-sx} dx \\
&= \int_0^{\infty} e^{(2-s)x} dx
\end{aligned}$$

To find the Laplace transform we need to compute the integral, the result of which will depend on the complex variable  $s$ . The Laplace transform is only defined when the real part of  $s$  is greater than the real part of the poles of the function (That's  $\Re(s) > 2$ ).

The integral is given by:

$$\begin{aligned}
\mathcal{L}\{f(x)\} &= \lim_{a \rightarrow \infty} \left[ \frac{e^{(2-s)x}}{2-s} \right]_0^a \\
&= \lim_{a \rightarrow \infty} \frac{e^{(2-s)a} - 1}{2-s}
\end{aligned}$$

if  $\Re(s) > 2$ ,  $2 - \Re(s)$  is negative and as  $a$  approaches infinity, the exponential term  $e^{(2-s)a}$  goes to 0, the limit becomes

$$\mathcal{L}\{f(x)\} = \lim_{a \rightarrow \infty} \frac{-1}{2-s} = \frac{1}{s-2}$$

(4)

We first need to express  $F(s)$  in partial fraction form and then find the inverse transforms of each term:

$$F(s) = \frac{1}{s^2 + 4s + 5} = \frac{1}{(s+2)^2 + 1}$$

Now, we can use the Laplace transform pair to find the inverse Laplace transform:

$$\mathcal{L}^{-1}\{e^{-ax} \sin(bx)\} = \frac{b}{(s+a)^2 + b^2}$$

That's  $a = 2, b = 1$ :

$$\mathcal{L}^{-1}\{f(x)\} = e^{-ax} \sin(bx) = e^{-2x} \sin(x)$$

### III. Numerical Methods

(1) Write a function that takes a list as input and returns the difference between each element and its succeeding element in the list.

(2) Write a function that takes a tuple representing an interval and solves the ordinary differential equation  $\frac{dy}{dx} = x - y$  with a step size of  $h = 0.1$ .

- Errata: In addition, the function also needs to accept a tuple representing the initial value

### Solutions (Haskell)

(1)

I've received some inquiries from readers regarding this question. They thought it involved providing a function expression for analysis and then receiving a set of discrete  $x$  inputs to calculate the differences. In reality, my intention is simply to calculate the differences between adjacent elements in a sequence.

```

forwardDifference :: Num a => [a] -> [a]
forwardDifference [] = []
forwardDifference [_] = []
forwardDifference (x:y:xs) = (y - x) : forwardDifference (y:xs)

```

Example for Test:

```

main :: IO ()
main = do
  let inputList = [1, 4, 7, 11, 16]
      diffList = forwardDifference inputList
  putStrLn $ "Input List: " ++ show inputList
  putStrLn $ "Forward Differences: " ++ show diffList

```

Output: [3,3,4,5]

(2)

We define a function representing the right-hand side of the ordinary differential equation, and within the Euler method function, we take an initial condition  $(x, y)$ , a solution interval  $(a, b)$ , and a step size  $h$ . We use the iterate function to construct an infinite list, where each element is the result of applying the Euler method to obtain the next point.

```

equation :: Double -> Double -> Double
equation x y = x - y

eulerMethod :: (Double -> Double -> Double)
             -> (Double, Double)
             -> (Double, Double)
             -> Double
             -> [(Double, Double)]
eulerMethod equation initialCondition interval stepSize = iterate step initial-
Condition
  where
    step (x, y) = (x + stepSize, y + stepSize * equation x y)

```

Example for Test:

```
main :: IO ()
main = do
  let initialCondition = (0, 1)
      interval = (0, 2)
      stepSize = 0.1

      solution = takeWhile (\(x, _) -> x <= snd interval)
                    $ eulerMethod equation initialCondition interval stepSize
      result = head $ dropWhile (\(x, _) -> x < 0.5) solution

  putStrLn $ "Result at x = " ++ show (fst result) ++ ", y = " ++ show (snd
result)
```

Output:

```
Result at x = 0.5: x = 0.5, y = 0.68098
```

(3)

This question is basically the same as the previous one, just change the formula and calculate it a few more times.

```

equation :: Double -> Double -> Double
equation x y = x + y

rungeKuttaMethod :: (Double -> Double -> Double)
                 -> (Double, Double)
                 -> (Double, Double)
                 -> Double
                 -> [(Double, Double)]
rungeKuttaMethod equation initialCondition interval stepSize = iterate step ini-
tialCondition
  where
    step (x, y) = let
      k1 = stepSize * equation x y
      k2 = stepSize * equation (x + 0.5 * stepSize) (y + 0.5 * k1)
      k3 = stepSize * equation (x + 0.5 * stepSize) (y + 0.5 * k2)
      k4 = stepSize * equation (x + stepSize) (y + k3)
    in (x + stepSize, y + (k1 + 2*k2 + 2*k3 + k4) / 6)

```

Example for Test:

```

main :: IO ()
main = do
  let equation x y = x + y
      initialCondition = (0, 1)
      interval = (0, 2)
      stepSize = 0.1

      solution = takeWhile (\(x, _) -> x <= snd interval)
                    $ rungeKuttaMethod equation initialCondition interval step-
Size
      result = head $ dropWhile (\(x, _) -> x < 0.5) solution

      putStrLn $ "Result at x = " ++ show (fst result) ++ ", y = " ++ show (snd
result)

```

Output:

```

Result at x = 0.5: x = 0.5, y = 1.7974412771936765

```